2,4-D	GROUP	4	HERBICIDE
Fluroxypyr	GROUP	4	HERBICIDE
Dicamba	GROUP	4	HERBICIDE

15-0-5

Turf King Fertilizer With Escalade® Weed & Feed

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

CONTAINS THE ESTERS OF 2,4-D AND FLUROXYPYR WITH DICAMBA ACID
Weeds contacted by this product gradually die and disappear from the lawn.
Treats (or covers) 12,500 Square Feet (100' x 125')

ACTIVE INGREDIENT:

Isooctyl (2-ethylhexyl) Ester of 2,4-Dichlorophenoxyacetic Acid*	0.8650%
1-Methylheptyl Ester of Fluroxypyr:((4-amino-3-5-dichloro-6-fluoro-2-pyridinyl)oxy)acetic Acid, 1-methylheptyl Ester **	0.2060%
Dicamba (3,6-Dichloro-o-Anisic Acid)***	0.0717%
OTHER INGREDIENTS:	98.8573%
TOTAL:	100.0000%

Isomer Specific Method, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid.	0.5740%
**((4-amino-3-5-dichloro-6-fluoro-2-pyridinyl)oxy)acetic Acid	0.1430%
***3,6-Dichloro-o-Anisic Acid	0.0717%

Fertilizer guaranteed by Andre and Son, Inc.

DO NOT SELL, DISTRIBUTE OR USE THE PRODUCT IN NASSAU AND SUFFOLK COUNTY IN NEW YORK.

KEEP OUT OF REACH OF CHILDREN

CAUTION

For Medical Emergencies, Call (877) 325-1840
For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing.

FIRST AID

IF IN EYES:
- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING:
- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 to 20 minutes.
- Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All loaders, applicators, and other handlers must wear:
- long-sleeved shirt and long pants,
- shoes plus socks,
- chemical-resistant gloves (Barrier Laminate, Butyl Rubber > 14 mils, Nitrile Rubber >14 mils, Neoprene rubber > 14 mils, or Viton> 14 mils)

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users Should:
- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Distributed and Guaranteed by Andre & Son
17150 State Route 706 • Montrose, PA 18801 • 570-278-1131

GUARANTEED ANALYSIS

Total Nitrogen (N)	15.00%
10.5% Urea Nitrogen	
4.5% CSRUN	
Soluble Potash (K2O)	5.00%

Nutrients derived from:
PCSCU (XCU), Urea, Muriate of Potash..

If this product is being sold in a state requiring the publication of information regarding its metal content, information regarding the contents and levels of metals in this product is available at the following website: http://www.aapfco.org/metals.htm

ENVIRONMENTAL HAZARDS

This pesticide may be toxic to fish and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

NON-TARGET ORGANISM ADVISORY STATEMENT: This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not enter or allow people (or pets) to enter the treated area until dusts have settled. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

This product is a stable mixture of three herbicides. It is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees and Roughs) and similar turf areas.

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as:

IMPORTANT RESTRICTIONS

This product is persistent and may be present in plant materials for over 30 days after application. Do not use treated plant material or manure from animals that have grazed or consumed forage from treated areas for compost, mulch, or mushroom spawn until 30 days after application.

Animals that have been fed fluroxypyr treated forage must be fed forage free of fluroxypyr for at least 3 days before they are moved off the treated property.

WEED RESISTANCE MANAGEMENT

For resistance management, this product is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to this product and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same area. Appropriate resistance management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of this product or other Group 4 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g. higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method. Prevent movement of resistant weed seeds to other areas by cleaning equipment.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Nufarm at 855-280-6609.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as:

Alder
Annual yellow sweet clover
Artichoke
Aster
Austrian fieldcress
Bedstraw
Beggartick
Biden
Bindweed
Bird vetch
Bitterweed
Bitter wintercress
Black-eyed Susan
Black medic
Black mustard
Black-seed plantain
Blessed thistle
Blue lettuce
Blue vervain
Box elder
Bracted plantain
Brassbuttons
Bristly oxtongue
Broadleaf dock
Broadleaf plantain
Broomweed
Buckhorn
Buckhorn plantain
Bulbous buttercup
Bull nettle
Bull thistle
Burdock
Burning nettle
Bur ragweed
Burweed
Buttercup
Canada thistle
Carolina geranium
Carpetweed
Catchweed bedstraw
Catsear
Catnip
Chickweed
Chicory
Cinquefoil
Clover
Cockle
Cocklebur
Coffeebean
Coffeeweed
Common chickweed
Common mullein
Common sowthistle
Corn Chamomile
Creeping jenny
Crimson clover
Croton
Cudweed
Curly dock
Curly indigo
Dandelion
Dead nettle
Dock
Dogbane
Dogfennel
Elderberry
English daisy
Fall dandelion
False dandelion
False flax
False sunflower
Fiddleneck
Field bindweed
Field pansy
Flea bane (daisy)
Flixweed
Florida betony
Florida pusley
Frenchweed
Galinsoga
Garlic mustard
Garlic mustard
Goathead
Goatsbeard
Goldenrod
Ground ivy
Gumweed
Hairy bittercress
Hairy fleabane
Hawkweed
Healall
Heartleaf drymary
Hedge bindweed
Hedge mustard
Hemp
Henbit
Hoary cress
Hoary plantain
Hoary vervain
Honeysuckle
Hop clover
Horsenettle
Horsetail
Indiana mallow
Ironweed
Jewelweed
Jimsonweed
Kochia
Knawel
Knotweed
Lambsquarter
Lespedeza
Locoweed
Lupine
Mallow
Marshelder
Matchweed
Mexicanweed
Milk vetch
Milkweed bloodflower
Mugwort
Morningglory
Mousear chickweed
Musk thistle
Mustard
Narrowleaf plantain
Narrowleaf vetch
Nettle
Orange hawkweed
Oxalis
Oxeye daisy
Parsley-piert
Parsnip
Pearlwort
Pennycress
Pennywort
Peppergrass
Pepperweed
Pigweed
Pineywoods bedstraw
Plains coreopsis
Plantain
Poison hemlock
Poison ivy
Poison oak
Pokeweed
Poorjoe
Povertyweed
Prickly lettuce
Prickly sida
Primrose
Prostrate knotweed
Prostrate pigweed
Prostrate spurge
Prostrate vervain
Puncture vine
Purslane
Ragweed
Red clover
Redroot pigweed
Red sorrel
Redstem filaree
Rough cinquefoil
Rough fleabane
Roundleafed marigold
Rush
Russian pigweed
Russian thistle
St. Johnswort
Scarlet pimpernel
Scotch thistle
Sheep sorrel
Shepherdspurse
Slender plantain
Smallflower galinsoga
Smartweed
Smooth dock
Smooth pigweed
Sneezeweed
Southern wild rose
Sowthistle
Spanishneedle
Spatterdock
Speedwell
Spiny Amaranth
Spiny cocklebur
Spotted catsear
Spotted knapweed
Spotted spurge
Spurge
Spurweed
Stinging nettle
Stinkweed
Stitchwort
Strawberry clover
Sumac
Sunflower
Sweet clover
Tall nettle
Tall vervain
Tansy mustard
Tansy ragwort
Tanweed
Tarweed
Thistle
Tick trefoil
Toadflax
Trailing Crownvetch
Tumble mustard
Tumble pigweed
Tumbleweed
Velvetleaf
Venice mallow
Veronica
Vervain
Vetch
Virginia buttonweed
Virginia creeper
Virginia pepperweed
Wavyleaf bullthistle
Western clematis
Western salsify
White clover
White mustard
Wild mustard
Wild aster
Wild buckwheat
Wild carrot
Wild four-o'-clock
Wild garlic
Wild geranium
Wild lettuce
Wild marigold
Wild onion
Wild parsnip
Wild radish
Wild rape
Wild strawberry
Wild sweet potato
Wild vetch
Willow
Witchweed
Wooly morningglory
Woodsorrel
Wooly croton
Wooly plantain
Wormseed
Yarrow
Yellow rocket
Yellowflower pepperweed
and other broadleaf weeds.

NET WT 50 lb (22.7 kg)

EPA Reg. No. 228-427-47956 • EPA Est. No. 47956-PA-001

DIRECTIONS FOR USE (continued)

WHEN TO APPLY: Best results are obtained when weeds are growing actively in spring or early fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain, sprinkling or dew); (3) Avoid unnecessary disturbances, including water of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed specified rates because damage to turf may occur.

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustinegrasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 50 pound bag should be used on 12,500 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 4 pounds per 1,000 square feet. These settings are furnished only as a guide. Age, spreader condition, operating speed, and unevenness of terrain can cause variation in applied rates. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound per 250 square feet.

SPREADER SETTINGS
ONE PASS APPLICATION*:

MAKE	SETTING
Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	4

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

Limitations: The maximum rate per application is 6 pounds per 1,000 square feet, limited to 2 applications per year. The maximum seasonal rate is 12 pounds per 1,000 square feet.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

ENVIRONMENTAL BENEFITS

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.

- 50' x 50' lawn produces enough oygen for a family of 4 every day
- Reduces Glare & Reduces Noise
- Traps Dust, Smoke Particles & Other Pollutants
- Purifies Water & Modifies Temperature
- Improves Soil Condition & Controls Erosion

Lot #

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not stack more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, or offered to others for their use, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

For nonrefillable bags of granular or dry formulation

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment, then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

or

Solid dilutable formulations in nonrefillable plastic containers

CONTAINER DISPOSAL: Nonrefillable Containers 5 Gallons or Less: Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

CONTAINER DISPOSAL: Nonrefillable containers larger than 5 Gallons: Nonrefillable container. Do not reuse or refill this container. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAkE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY kIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

LIMITATION OF LIABILITY

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR RISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded. (RV110526)

Escalade is a Registered Trademark of Nufarm Americas Inc. All other trademarks are the property of their respective owners.