

ACTIVE INCREDIENTS

COMMERCIAL

# **Turf King Fertilizer** With Viper® Weed & Feed

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

CONTAINS 2.4-D. MECOPROP-p AND DICAMBA

Weeds contacted by this product gradually die and disappear from the lawn.

Treats (or covers) 12,500 Square Feet (125' x 100')

#### **GET THE OPTICAL ADVANTAGE®**

ACTIVE INGREDIENTS:	
Isooctyl (2-ethylhexyl) Ester of 2,4-Dichlorophenoxyacetic Acid*	
2,4-Dichlorophenoxyacetic Acid*	0.146%
(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid**‡	0.287%
Dicamba (3,6-Dichloro-o-Anisic Acid)***	
OTHER INGREDIENTS:	98.420%
TOTAL:	100.000%
Isomer Specific Method, Equivalent to:	
*2,4-Dichlorophenoxyacetic Acid (Total)	0.860%
**(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid	0.287%
***3,6-Dichloro-o-Anisic Acid	0.070%
‡CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-p	

Fertilizer guaranteed by Andre and Son, Inc.

## **KEEP OUT OF REACH OF CHILDREN CAUTION**

SEE BACK PANEL FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300 For Medical Emergencies Only, Call (877) 325-1840

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are any waterproof materials. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

#### All mixers, loaders, applicators, flaggers, and other handlers must wear:

- · long-sleeved shirt and long pants,
- · shoes and socks, plus
- · chemical-resistant gloves (except for applicators using groundboom equipment)

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### **USER SAFETY RECOMMENDATIONS**

## Users Should:

- · Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- · Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## **FIRST AID**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF IN EYES:

- · Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- · Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- · Call a poison control center or doctor for treatment advice.

#### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment You may also contact 1-877-325-1840 for emergency medical treatment information.

> Distributed and Guaranteed by Andre & Son 17150 State Route 706 • Montrose, PA 18801 • 570-278-1131

#### **GUARANTEED ANALYSIS**

12.0% Urea Nitrogen 3.0% CSRUN 

#### Nutrients derived from:

XCU, Urea, Muriate of Potash.

If this product is being sold in a state requiring the publication of information regarding its metal content, information regarding the contents and levels of metals in this product is available at the following website: http://www.aapfco.org/metals.htm

#### **ENVIRONMENTAL HAZARDS**

This pesticide may be toxic to fish and aquatic invertebrates and may be adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater

## **DIRECTION FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling, READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not enter or allow people (or pets) to enter the treated area until dusts have settled. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Do not use this product on or near desirable plants, including within the dripline of the roots of desirable trees and shrubs, since injury may result.

This product is a stable mixture of four herbicides. It is for use on Ornamental Turf lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees and Roughs) and similar turf areas

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth in lawns and controls many broadleaf weeds such as:

Alder	Box elder	Catchweed bedstraw	Curly dock
Annual yellow	Bracted plantain	Catsear	Curly indigo
sweet clover	Brassbuttons	Catnip	Dandelion
Artichoke	Bristly oxtongue	Chickweed	Dead nettle
Aster	Broadleaf dock	Chicory	Dock
Austrian fieldcress	Broadleaf plantain	Cinquefoil	Dogbane
Bedstraw	Broomweed	Clover	Dogfennel
Beggartick	Buckhorn	Cockle	Elderberry
Biden	Buckhorn plantain	Cocklebur	English daisy
Bindweed	Bulbous buttercup	Coffeebean	Fall dandelion
Bird vetch	Bull nettle	Coffeeweed	False dandelion
Bitterweed	Bull thistle	Common chickweed	False flax
Bitter wintercress	Burdock	Common mullein	False sunflower
Black-eyed Susan	Burning nettle	Common sowthistle	Fiddleneck
Black medic	Bur ragweed	Corn	Field bindweed
Black mustard	Burweed	Chamomile	Field pansy
Blackseed plantain	Buttercup	Creeping jenny	Flea bane (daisy)
Blessed thistle	Canada thistle	Crimson clover	Flixweed
Blue lettuce	Carolina geranium	Croton	Florida betony
Blue vervain	Carpetweed	Cudweed	Florida pusley

NET WT 50 lb (22.7 kg)

Frenchweed Nettle Galinsoga Orange hawkweed Garlic mustard Oxalis Goathead Oxeye daisy Goatsbeard Parslev-piert Goldenrod Parsnip Pearlwort Ground ivv Gumweed Pennycress Hairy bittercress Pennywort Hairy fleabane Peppergrass Hawkweed Pepperweed Healall Pigweed Heartleaf dryman Pinevwoods bedstraw Hedge bindweed Plains coreopsis Hedge mustard Plantain Musk thistle Hemp Henbit Mustard Hoary cress Narrowleaf plantain Narrowleaf vetch Hoary plantain Hoary vervain Nettle Honeysuckle Orange hawkweed Hop clover Oxalis Horsenettle Oxeye daisy Parsley-piert Horsetail Indiana mallow Parsnip Pearlwort Ironweed Jewelweed Pennycress .limsonweed Pennywort Peppergrass Kochia Knawel Pepperweed Knotweed Pigweed Pineywoods bedstraw Lambsquarter Lespedeza Plains coreopsis Plantain Locoweed Lupine Poison hemlock Mallow Poison ivy Marshelder Poison oak Matchweed Pokeweed Mexicanweed Poorjoe Milk vetch Povertyweed Prickly lettuce Milkweed bloodflower

Prickly sida

Prostrate knotweed

Prostrate pigweed

Prostrate spurge

Prostrate vervain

Puncture vine

Primrose

Mugwort

Mustard

Morningglory

Musk thistle

Mousear chickweed

Narrowleaf plantain

Narrowleaf vetch

Ragweed Red clover Redroot pigweed Red sorrel Redstem filaree Rough cinquefoil Rough fleabane Roundleafed marigold Rush Russian pigweed Russian thistle St .lohnswort Scarlet pimpernel Scotch thistle Sheep sorrel Shepherdspurse Slender plantain Smallflower galinsoga Smartweed Smooth dock Smooth pigweed Sneezeweed Southern wild rose Sowthistle Spanishpeedle Spatterdock Speedwell Spiny Amaranth Spiny cocklebur Spotted catsear Spotted knapweed Spotted spurge Spurge Spurweed Stinging nettle Stinkweed Stitchwort Strawberry clover Sumac Sunflower Sweet clover Tall nettle Tall vervain Tansv mustard Tansy ragwort

Purslane

Tumbleweed Velvet leaf Venice mallow Veronica Vervain Vetch Virginia buttonweed Virginia creeper Virginia pepperweed Wavvleaf bullthistle Western clematis Western salsify White clover White mustard Wild mustard Wild aster Wild buckwheat Wild carrot Wild four-o'-clock Wild garlic Wild geranium Wild lettuce Wild marigold Wild onion Wild parsnip Wild radish Wild rape Wild strawberry Wild sweet potato Wild vetch Willow Witchweed Woolly morningglory Woodsorrel Woolly croton Woolly plantain Wormseed Yarrow Yellow rocket Yellowflower Tanweed pepperweed Tarweed

Thistle

Tick trefoil

Toadflax Trailing Crownvetch

Tumble mustard

Tumble pigweed

#### WHEN TO APPLY

Best results are obtained when weeds are growing actively in spring or early fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers; especially when applying with a rotary (spinner) type spreader. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. For optimum results (1) Avoid mowing 1 to 2 days before and after application; (2) Apply when grass is moist (after a rain sprinkling or dew); (3) Avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material; and (4) Avoid applying if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

#### **HOW TO APPLY**

This 50 pound bag should be used on 12,500 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is equal to one application.

The following settings are approximate for applying contents of this package at a suggested rate of 4 pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound per 250 square

EPA Reg. No. 228-413-47956 • EPA Est. No. 47956-PA-001

## **DIRECTION FOR USE (continued)**

#### SPREADER SETTINGS ONE PASS APPLICATION<sup>°</sup>:

MAKE	SETTING
Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	4

Sweep (do not wash) all fertilizer materials from surfaces immediately after application, because certain fertilizer ingredients can cause permanent stains. Wash out spreader after each use and allow it to dry thoroughly. Treat any exposed metal spreader areas with oil to reduce corrosion.

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing \*(No. of lbs.) pounds of nitrogen per 1,000 square feet (optional  $P_2O5$  and  $K_2O$  levels) and herbicide to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

\*Based on Inert fertilizer claimed

#### Limitations

The maximum rate per application is 4 pounds per 1,000 square feet, limited to 2 applications per year. The maximum seasonal rate is 8 pounds per 1,000 square feet. Minimum of 30 days between applications.

#### **ENVIRONMENTAL BENEFITS**

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.

- A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- Reduces Glare & Reduces Noise
- Traps Dust, Smoke Particles & Other Pollutants
- Purifies Water & Modifies Temperature
- Improves Soil Condition & Controls Erosion

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not stack more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment, then offer for recycling if available, or dispose of empty bag in a sanitary landfill or by incineration or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

#### WARRANTY DISCLAIMER

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PRAPE USAGE, UNINTED DESAGE. UNINTED DESAGES, INCLUDING BUT NOT LIMITED TO INFEFRED TWENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

### **LIMITATION OF LIABILITY**

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. TO THE EXTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BY USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR RISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.

If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.

Viper and The Optical Advantage are Registered Trademarks of Nufarm Americas Inc.

(RV061417)